

AD-A070 929

TEXAS UNIV AT AUSTIN DEPT OF CHEMISTRY

F/6 7/4

A UV PHOTOELECTRON SPECTROSCOPIC INVESTIGATION OF THE BONDING I--ETC(U)

1979 M H CHISHOLM, A H COWLEY, M LATTMAN

N00014-76-C-0577

UNCLASSIFIED

TR-79-01

NL

OF  
AD  
A070 929



UNCLASSIFIED

LEVEL 1

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

## REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS  
BEFORE COMPLETING FORM

1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A UV PHOTOELECTRON SPECTROSCOPIC INVESTIGATION OF THE BONDING IN SOME TRI-, TETRA-, AND PENTACOORDINATED DIALKYLAMINO COMPOUNDS OF CHROMIUM, MOLYBDENUM, NIOBIUM, AND TANTALUM		5. TYPE OF REPORT & PERIOD COVERED Technical Report, 1979
7. AUTHOR(s) M. H. Chisholm*, A. H. Cowley*, and M. Lattman		6. PERFORMING ORG. REPORT NUMBER TR 79-01
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry The University of Texas at Austin Austin, Texas 78712		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0577
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) See page 1.		12. REPORT DATE 1979
		13. NUMBER OF PAGES 23
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release: Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DDC RECEIVED JUL 9 1979 D		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) UV Photoelectron Spectroscopic, Dialkylamino, Bonding, Chromium, Molybdenum, Niobium, Tantalum		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Helium (I) ultraviolet photoelectron spectra (UPS) are reported for the compounds $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ , $\text{Cr}(\text{NET}_2)_4$ , $\text{Mo}(\text{NME}_2)_4$ , $\text{Mo}(\text{NET}_2)_4$ , $\text{Nb}(\text{NME}_2)_5$ , and $\text{Ta}(\text{NME}_2)_5$ . The interpretation of the UPS of the paramagnetic dialkylamide, $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ , was aided by a $\text{SCF-X}_\alpha$ scattered wave ( $\text{X}_\alpha\text{-SW}$ ) calculation on the model $\rightarrow$		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

A070929

DDC FILE COPY

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

→ compound  $\text{Cr}(\text{NH}_2)_3$ . In contrast to previous UPS work on  $\text{Cr}[\text{N}(\text{SiMe}_3)_2]_3$ , ionizations are detectable from the metal MO's of  $\text{Cr}(\text{N}^1\text{Pr}_2)_3$ . The UPS of the tetracoordinate molybdenum compounds,  $\text{Mo}(\text{NR}_2)_4$ ,  $\text{R}=\text{Me}$ ,  $\text{Et}$  are in excellent agreement with X-ray crystallographic data on  $\text{Mo}(\text{NMe}_2)_4$  and exhibit a low energy ionization ( $\sim 5.3$  eV) which is attributable to electron ejection from the essentially pure  $\text{Mo } 4d_{(x)}^2(-y)^2$  MO of  $b_1$  symmetry. The pentacoordinate compounds,  $\text{M}(\text{NMe}_2)_5$ ,  $\text{M}=\text{Nb}$ ,  $\text{Ta}$  exhibit very similar UPS and therefore appear to be isostructural in the vapor phase. ↙

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist.	Avail and/or special
A	

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

6

A UV PHOTOELECTRON SPECTROSCOPIC INVESTIGATION  
OF THE BONDING IN SOME TRI-, TETRA-, AND  
PENTACOORDINATED DIALKYLAMINO COMPOUNDS OF  
CHROMIUM, MOLYBDENUM, NIOBIUM, AND TANTALUM.

12 28 p.

10

M. H. / Chisholm<sup>7a\*</sup>, A. H. / Cowley<sup>7b\*</sup> and M. / Lattman<sup>7c</sup>

14

TR-79-01

15

NO 79-76-C-0577

Contribution from the ~~Departments of Chemistry~~,  
 Indiana University, Bloomington, Indiana and  
 the University of Texas at Austin, Austin, Texas 78712  
 Received \_\_\_\_\_

9

Technical Repts.

11

1979

Abstract: Helium (I) ultraviolet photoelectron spectra (UPS) are reported for the compounds  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ ,  $\text{Cr}(\text{NEt}_2)_4$ ,  $\text{Mo}(\text{NMe}_2)_4$ ,  $\text{Mo}(\text{NEt}_2)_4$ ,  $\text{Nb}(\text{NMe}_2)_5$ , and  $\text{Ta}(\text{NMe}_2)_5$ . The interpretation of the UPS of the paramagnetic dialkylamide,  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ , was aided by a SCF  $X_\alpha$  scattered wave ( $X_\alpha$  SW) calculation on the model compound  $\text{Cr}(\text{NH}_2)_3$ . In contrast to previous UPS work on  $\text{Cr}[\text{N}(\text{SiMe}_3)_2]_3$ , ionizations are detectable from the metal MO's of  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ . The UPS of the tetracoordinate molybdenum compounds,  $\text{Mo}(\text{NR}_2)_4$ ,  $\text{R} = \text{Me}, \text{Et}$  are in excellent agreement with X-ray crystallographic data on  $\text{Mo}(\text{NMe}_2)_4$  and exhibit a low energy ionization ( $\sim 5.3$  eV) which is attributable to electron ejection from the essentially pure  $\text{Mo } 4d_{x^2-y^2}$  MO of  $b_1$  symmetry. The pentacoordinate compounds,

347 830

sep

$M(NMe_2)_5$ ,  $M = Nb, Ta$  exhibit very similar UPS and therefore appear to be isostructural in the vapor phase.

## INTRODUCTION

The use of dialkylamido and disilylamido ligands has permitted the isolation of mono- and dinuclear transition metal derivatives with widely varying coordination numbers.<sup>2</sup> Compounds of this type are well suited to investigation by ultraviolet photoelectron spectroscopy (UPS) because, as shown in previous studies<sup>3-5</sup>, the peaks at lower binding energies corresponding to electron ejection from metal-centered MO's, nitrogen lone pair MO's, and metal-nitrogen  $\sigma$ -bonding MO's fall into distinct regions. The UPS data can be used, therefore, not only for the delineation of bonding schemes but also for gaining insights into the stereochemistry of metal amides.

The present paper is concerned with the measurement and interpretation of the UPS of  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ ,  $\text{Cr}(\text{NEt}_2)_4$ ,  $\text{Mo}(\text{NMe}_2)_4$ ,  $\text{Mo}(\text{NEt}_2)_4$ ,  $\text{Nb}(\text{NMe}_2)_5$  and  $\text{Ta}(\text{NMe}_2)_5$ . Our interest in the tricoordinate species,  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ , was generated by the fact that in the corresponding silylamide,  $\text{Cr}[\text{N}(\text{SiMe}_3)_2]_3$ , it was impossible to detect UPS peaks corresponding to the ionization of metal-localized MO's.<sup>5</sup> The tetracoordinate dialkylamides,  $\text{Cr}(\text{NEt}_2)_4$  and  $\text{Mo}(\text{NR}_2)_4$ ,  $\text{R} = \text{Me}, \text{Et}$ , are the first Group VI A amides to be studied by UPS; particular interest is associated with these compounds because the molybdenum amides are diamagnetic yet  $\text{Cr}(\text{NEt}_2)_4$  is paramagnetic. Finally, the pentacoordinate amides,  $\text{M}(\text{NMe}_2)_5$ ,  $\text{M} = \text{Nb}, \text{Ta}$ , were investigated because of the apparently close energies of the trigonal bipyramidal and square pyramidal  $\text{MN}_5$  geometries and the possibility

of detecting conformational effects in the vapor phase.

A SCF  $X_\alpha$  scattered-wave calculation<sup>6</sup> (hereafter referred to as  $X_\alpha$  SW) has been performed on the model amide  $\text{Cr}(\text{NH}_2)_3$  to facilitate the interpretation of the UPS data. As is well known, this theoretical approach has the advantage of avoiding the use of Koopmans' theorem<sup>7</sup> by means of the transition state method<sup>8</sup>.

## EXPERIMENTAL SECTION

Materials. The compounds  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ <sup>9</sup>,  $\text{Cr}(\text{NEt}_2)_4$ <sup>10</sup>,  $\text{Mo}(\text{NMe}_2)_4$ <sup>11</sup>,  $\text{Mo}(\text{NEt}_2)_4$ <sup>11</sup>,  $\text{Nb}(\text{NMe}_2)_5$ <sup>12</sup>, and  $\text{Ta}(\text{NMe}_2)_5$ <sup>13</sup> were prepared and purified as described in the literature.

Spectroscopic Measurements. All UPS data were recorded on a Perkin-Elmer Model PS-18 Photoelectron Spectrometer using a He(I) photon source (21.22 eV). The heated inlet probe was used for all samples and temperatures in the range 35–100°C were necessary to obtain suitable spectra. Each spectrum was calibrated with xenon (12.130 eV) and argon (15.759 eV) used as internal standards. Spectral resolution was maintained between 25 and 50 meV for the argon line. All ionization energies are read as the band maxima, unless otherwise noted, and are the average of at least three different runs.


Computational Procedures. The  $X_\alpha$  SW calculation on  $\text{Cr}(\text{NH}_2)_3$  was made by employing the spin-restricted procedure of Johnson and Slater<sup>6</sup>, setting the occupation numbers of the 6e and 4a<sub>1</sub> levels at two and one, respectively, in accord with the

observed ground state electronic configuration.<sup>9</sup> The bond distances and angles for  $\text{Cr}(\text{NH}_2)_3$  were based on those observed by X-ray crystallography for  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ ,<sup>14</sup> except the N-H bond distance was taken to be 1.01 Å. The atomic sphere radii were chosen on the basis of optimizing the virial ratio, and the outer sphere was set tangential to the hydrogen spheres.<sup>15</sup> Schwartz's exchange parameters,  $\alpha_{\text{HF}}$ , were used for chromium and nitrogen, while Slater's value for hydrogen was used.<sup>16</sup>  $\alpha_{\text{OUTER}}$  was taken equal to  $\alpha_{\text{H}}$ , while the intersphere exchange parameter,  $\alpha_{\text{INT}}$ , was calculated to be 0.7490 on the basis of averaging the atomic  $\alpha$  values according to the numbers of valence electrons. Spherical harmonics through  $\ell = 2$  were employed for the chromium and outer spheres, while functions through  $\ell=1$  and  $\ell=0$  were employed for the nitrogen and hydrogen spheres, respectively. All SCF calculations were converged to better than 0.01 eV for each level, maintaining all cores fixed. The first four IE's for  $\text{Cr}(\text{NH}_2)_3$  were computed by the transition state method<sup>8</sup>.

## RESULTS AND DISCUSSION

The tricoordinate compounds,  $\text{M}[\text{N}(\text{SiMe}_3)_2]_3$ ,  $\text{M}=\text{Ti}$ ,  $\text{Cr}$ ,  $\text{Fe}$ , and  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$  represent examples of the stabilization of low coordination numbers by means of bulky groups. Single

crystal X-ray structures of  $\text{Fe}[\text{N}(\text{SiMe}_3)_2]_3$ <sup>17</sup> and  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ <sup>14</sup> revealed that the  $\text{MN}_3$  skeletons are trigonal planar and that  $\phi$ , the dihedral angle between the  $\text{NX}_2$  and  $\text{MN}_3$  planes, is  $49^\circ$  and  $71^\circ$ , respectively. The other tricoordinate compounds of concern here can be assumed to possess a similar structure. Collectively, ESR data<sup>14,18</sup>, magnetic data<sup>19</sup>, and crystal field calculations<sup>19</sup> indicate that the ground state electronic configurations of the silylamides are



The dialkylamide,  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ , has been shown<sup>9</sup> to have the same ground state electronic configuration as  $\text{Cr}[\text{N}(\text{SiMe}_3)_2]_3$ .

Curiously, despite the presence of unpaired electrons in the above silylamides no low energy ionizations attributable to metal-localized orbitals were detectable in the UPS<sup>5</sup>.

The reason which was advanced for this phenomenon is that the metal MO's are stabilized significantly by the more electron-withdrawing  $(\text{Me}_3\text{Si})_2\text{N}$  groups, possibly causing them to be of comparable energies to the nitrogen lone pair MO's. Interestingly,  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$  does exhibit peaks at low IE which can be attributed to the ionization of metal-localized MO's (Figure 1a). The detailed assignment of the UPS of paramagnetic transition metal systems is, in general, a complicated matter<sup>20,21</sup> because of the large number of ionic states which can be generated upon photoionization (Table I). Arguing qualitatively, and on the basis of intensity considerations it is reasonable to assign peaks  $I_1$  and  $I_2$  to the production of the  $^3A_2$  and  $^3E$  ionic states via electron ejection from the  $\text{Cr}(3d)$  orbitals of  $a_2$  and  $e$  symmetry, respectively (see below). In threefold symmetry the nitrogen "lone pair" MO's span the  $a_2$  and  $e$  irreducible representations. However, as is clear from the nodal properties of such MO's (Figure 2) their relative energies are sensitive to the dihedral angle.<sup>22</sup> Obviously, at intermediate values of  $\phi$ , a "cross-over" occurs, and, at some particular value of  $\phi$  the  $a_2$  and  $e$  MO's must be degenerate. Since the dihedral angle between the  $\text{CrN}_3$  and  $\text{NC}_2$  planes is  $71^\circ$  in  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ <sup>14</sup> the symmetry is nearer to  $D_{3h}(1)$  than  $D_{3h}(2)$ . It is on this basis that we assign peaks  $I_3$  and  $I_4$  to the ionization of the  $a_2$  and  $e$  nitrogen lone pair MO's respectively. The lack of resolution of peaks  $I_3$  and  $I_4$  could be due to two factors: (a) the possibility that both triplet and quintet  $A_2$  and  $E$  states are observed in this region and (b) Jahn-Teller splitting of the  $E$  states. The

peak at 9.9 eV obviously represents the onset of ionization of the metal-nitrogen  $\sigma$ -bonding MO's (of symmetry  $a_1$  and e); however, it is difficult to advance specific assignments.

The foregoing, qualitative interpretation of the UPS of  $\text{Cr}(\text{N}^1\text{Pr}_2)_3$  is in good accord with the  $X_\alpha$  SW calculation on the model compound  $\text{Cr}(\text{NH}_2)_3$  (Tables II and III). Thus, the two highest occupied MO's (Table II) are primarily Cr(3d) in composition; the 6e MO is the degenerate  $3d_{xz}, d_{yz}$  set while the  $4a_1$  is primarily  $3d_z^2$  (if the  $\text{CrN}_3$  moiety lies in the xy plane). The  $3a_2$  and 5e MO's are both mainly nitrogen lone pair; however, while the  $3a_2$  is entirely localized on the nitrogens, the 5e has a small, but significant amount of density on chromium. This orbital is the  $\pi$ -bonding MO of the molecule and the orbital contour of one of the degenerate set is shown (in the xy plane) in Figure 3a. Since the nitrogen lone pairs are twisted out of the xy plane by only  $19^\circ$ , the best overlap is with the  $d_{x^2-y^2}, d_{xy}$  set of orbitals on chromium. The wavefunction in Figure 3a shows nitrogen lone pair overlap with the Cr  $3d_{xy}$  orbital. The next two MO's, 4e and  $3a_1$ , are the  $\sigma$ -bonding orbitals of the molecule. The contour of one of the degenerate 4e wavefunctions is shown in Figure 3b (xy plane), while the contour of the  $3a_1$  MO (in the plane along the Cr-N  $\sigma$  bond, perpendicular to the xy plane) is illustrated in Figure 3c. The  $3d_z^2$  contribution to the  $\sigma$ -bonding is evidenced from the latter contour. The next four MO's,  $2a_2$ , 3e, 2e, and  $2a_1$ , are N-H  $\sigma$ -bonding, while the  $1a_2$ , 1e, and  $1a_1$ , are entirely Cr 3p and 3s in composition. It appears that the Cr  $3d_{x^2-y^2}, d_{xy}$  set of e orbitals is involved in  $\pi$ -bonding (Figure 3a) as well as  $\sigma$ -bonding (Figure 3b) in the molecule.

It is of importance to note that the computed IE's (Table III) for the  $4a_1$  and  $6e$  metal-localized MO's are in the reverse order to the ground state eigenvalues (Table II), i.e. the HOMO ( $6e$ ) is responsible for the second ionization. The breakdown of Koopmans' theorem in this instance is due to the relatively large relaxation energies which are associated with the photoionization of metal-rich orbitals: the higher the metal  $d$  orbital character, the more the MO will relax upon ionization. Thus, the  $6e$  (87% metal) relaxes to a larger extent than the  $4a_1$  (71% metal), causing the crossover in the observed UPS. This type of behavior is characteristic of the UPS of several transition metal compounds.<sup>23</sup>

The tetracoordinate dialkylamides of Cr and Mo are of interest because, while both are  $d^2$  systems,  $Cr(NEt_2)_4$  is paramagnetic yet  $Mo(NMe_2)_4$  and  $Mo(NEt_2)_4$  are diamagnetic. If the nitrogen geometry is taken to be trigonal planar and metal-nitrogen bond rotation is assumed to be slow on the UPS time scale, the maximum skeletal symmetry of a  $M(NR_2)_4$  compound is  $D_{2d}$ . There are, in fact, two possible structures that possess this geometry:  $D_{2d}(1)$  in which the C-N-C planes are perpendicular to the dihedral planes, and  $D_{2d}(2)$  in which the C-N-C

planes are parallel to the dihedral planes. In  $T_d$  symmetry the metal nd orbitals split into the familiar e and  $t_2$  sets; reduction of symmetry to  $D_{2d}(1)$  or  $D_{2d}(2)$  results in these orbitals transforming as  $b_1$ ,  $a_1$ ,  $b_2$ , and e as illustrated in Figure 4. In  $D_{2d}(1)$  symmetry the nitrogen lone pair MO's,  $n_N$ , span the irreducible representations  $a_1$ ,  $b_2$ , and e while in  $D_{2d}(2)$  symmetry these transform as  $a_2$ ,  $b_1$ , and e. Dative  $\pi$  bonding interactions are permitted between the nitrogen lone pair and metal nd MO's of the same symmetry as illustrated. Thus it is clear (Figure 4) that no interactions are possible with the  $b_1$  ( $d_{x^2-y^2}$ ) or  $a_1$  ( $d_z^2$ ) metal and MO's in  $D_{2d}(1)$  or  $D_{2d}(2)$  symmetry, respectively. These orbitals are the HOMO's in low spin  $d^2$  systems.

The crystal structure of  $Mo(NMe_2)_4$  has been investigated recently<sup>24</sup> and the molecule has been found to adopt  $D_{2d}(1)$  symmetry. The UPS of  $Mo(NMe_2)_4$  (Figure 1b) is consistent with the ground state electronic configuration... $(a_1)^2(b_2)^2(e)^4(b_1)^2$ . Thus,  $I_1$  clearly corresponds to ionization of the  $b_1$  metal localized MO, while peaks  $I_2$ ,  $I_3$ , and  $I_4$  can be assigned to electron ejection from the nitrogen lone pair MO's of symmetries e,  $b_2$ , and  $a_1$ , respectively on the basis of intensity considerations. The broad, intense peak at 10.7 eV represents ionization of the  $a_1$ , e, and  $b_2$  molybdenum-nitrogen  $\sigma$ -bonding MO's of symmetries  $a_1$ , e, and  $b_2$ . The general correctness of this sequence of orbitals has been confirmed by Fenske-Hall calculations on  $Mo(NMe_2)_4$  which, in addition, revealed that the  $b_1$  MO is 97%  $4d_{x^2-y^2}$  in character, and that the e,  $b_2$ , and  $a_1$  MO's

comprise 91, 94, and 71% ligand  $\pi$  character, respectively.<sup>24</sup> The low value for  $I_1$  (5.30 eV) is consistent with the Fenske-Hall calculations on  $\text{Mo}(\text{NMe}_2)_4$  which indicate that the  $b_1$  MO is slightly antibonding. The UPS of  $\text{Mo}(\text{NEt}_2)_4$  (Figure 1c) is quite similar to that of  $\text{Mo}(\text{NMe}_2)_4$  and can be assigned in an analogous manner. As expected the energy required for production of the  $^2B_1$  state via electron ejection from the  $\text{Mo}(4d_{x^2-y^2})$  orbital is virtually identical in both compounds, and slight inductive shifts are apparent in the IE's associated with the nitrogen lone pair MO's. The somewhat more complex appearance of the spectrum of  $\text{Mo}(\text{NEt}_2)_4$  in the nitrogen lone pair region could be due to (a) the actual symmetry of  $\text{Mo}(\text{NEt}_2)_4$  being lower than that of  $\text{Mo}(\text{NMe}_2)_4$ , or (b) an increased Jahn/Teller splitting of the  $^2E$  state in  $\text{Mo}(\text{NEt}_2)_4$  compared to  $\text{Mo}(\text{NMe}_2)_4$ .

Interestingly,  $\text{Cr}(\text{NEt}_2)_4$  (Figure 1d) is paramagnetic, although the reasons for this are not clear at the present time. Probably it is a reflection of the spin-pairing energies being in the order  $\text{Cr} > \text{Mo}$  and the ligand field energies being in the order  $\text{Mo} > \text{Cr}$ ; alternatively,  $\text{Cr}(\text{NEt}_2)_4$  could adopt the  $D_{2d}(2)$  structure. If so, the electronic configuration of the ground state would be  $(b_1)^2(e)^4(a_2)^2(a_1)^1(b_1)^1$  rather than  $(a_1)^2(b_2)^2(e)^4(b_1)^1(a_1)^1$ . The ionic states emerging from the photoionization of both the  $D_{2d}(1)$  and  $D_{2d}(2)$  ground state electronic configurations are presented in Table I along with possible spectral assignments. If one argues (probably in an oversimplified way) on the basis of peak intensities the better fit is obtained with the  $D_{2d}(2)$  ground state structure. However, this conjecture, while

interesting, should be viewed cautiously. Clearly, the structure of  $\text{Cr}(\text{NEt}_2)_4$  should be determined.

One of the intriguing aspects of the pentacoordinate amides concerns the apparently close energies of the square pyramidal and trigonal bipyramidal  $\text{MN}_5$  geometries. Thus, it has been demonstrated by X-ray crystallography that the  $\text{NbN}_5$  moiety of  $\text{Nb}(\text{NMe}_2)_5$  approaches a square pyramidal structure<sup>25</sup>, while  $\text{Ta}(\text{NEt}_2)_5$  has been found to adopt a trigonal bipyramidal geometry for the  $\text{TaN}_5$  skeleton.<sup>26</sup> The UPS of  $\text{Nb}(\text{NMe}_2)_5$  and  $\text{Ta}(\text{NMe}_2)_5$  are virtually identical (except for a small shift in the absolute value of the IE's) and only the spectrum of the latter is illustrated in Figure 1e. It thus appears that  $\text{Nb}(\text{NMe}_2)_5$  and  $\text{Ta}(\text{NMe}_2)_5$  are isostructural in the vapor phase. In a square pyramidal ( $\text{C}_2$ ) arrangement of  $\text{MNC}_2$  moieties the five peaks in the nitrogen lone pair region would correspond to the ionization of two a and three b MO's. However, it is difficult to be more specific without MO calculations. The lower IE's are, therefore, listed without assignments in Table IV.

**ACKNOWLEDGMENT,** The authors are grateful to the Office of Naval Research (Contract N00014-76-C-0577, Task No. NR 053-612) for financial support.

## REFERENCES AND NOTES

- (1) (a) Indiana University; (b) University of Texas at Austin.
- (2) For reviews, see (a) D. C. Bradley, *Adv. Inorg. Chem. Radiochem.*, **15**, 259 (1972); (b) D. C. Bradley and M. H. Chisholm, *Acc. Chem. Res.*, **9**, 273 (1976).
- (3) S. G. Gibbins, M. F. Lappert, J. B. Pedley, and G. J. Sharp, *J. Chem. Soc. Dalton Trans.*, 72 (1975).
- (4) D. H. Harris, M. F. Lappert, J. B. Pedley, and G. J. Sharp, *J. Chem. Soc. Dalton Trans.*, 945 (1976).
- (5) M. F. Lappert, J. B. Pedley, G. J. Sharp, and D. C. Bradley, *J. Chem. Soc. Dalton Trans.*, 1737 (1976).
- (6) J. C. Slater, *Adv. Quantum Chem.*, **6**, 1 (1972); (b) K. H. Johnson, *ibid.*, **7**, 143 (1973); (c) K. H. Johnson, *Annu. Rev. Phys. Chem.*, **26**, 39 (1975).
- (7) T. Koopmans, *Physica*, **1**, 104 (1934).
- (8) (a) J. C. Slater and K. H. Johnson, *Phys. Rev.*, **B**, **5**, 844 (1972); (b) J. C. Slater, Quantum Theory of Molecules and Solids, Vol. 4, McGraw-Hill, New York, 1974, p. 583.
- (9) E. C. Alyea, J. S. Basi, D. C. Bradley, and M. H. Chisholm, *Chem. Commun.*, 495 (1968).
- (10) J. S. Basi and D. C. Bradley, *Proc. Chem. Soc. Landau*, 305 (1963).
- (11) D. C. Bradley and M. H. Chisholm, *J. Chem. Soc. A*, 274 (1971).
- (12) D. C. Bradley and I. M. Thomas, *Can. J. Chem.*, **40**, 449 (1962).
- (13) D. C. Bradley and I. M. Thomas, *Can. J. Chem.*, **40**, 1355 (1962).

- (14) D. C. Bradley, M. B. Hursthouse, and C. W. Newing, *Chem. Commun.*, 411 (1971).
- (15) (a) J. G. Norman, *J. Chem. Phys.*, **61**, 4630 (1974);  
(b) J. G. Norman, *Mol. Phys.*, **31**, 1191 (1976). (c)  $r_{cr} = 2.586$ ,  $r_N = 1.136$ ,  $r_H = 0.7724$ ,  $r_{OUTER} = 5.537$  (in a.u.).
- (16) (a) K. Schwartz, *Phys. Rev.*, **B 5**, 2466 (1972). (b) J. C. Slater, *Int. J. Quantum Chem.*, **1**, 533 (1973).
- (17) D. C. Bradley, M. B. Hursthouse, and P. F. Rodesiler, *Chem. Commun.*, 14 (1969).
- (18) (a) D. C. Bradley and R. G. Copperthwaite, *Chem. Commun.*, 764 (1971). (b) E. C. Alyea, D. C. Bradley, R. G. Copperthwaite, K. D. Sales, B. W. Fitzsimmons, and C. E. Johnson, *Chem. Commun.*, 1715 (1970).
- (19) E. C. Alyea, D. C. Bradley, R. C. Copperthwaite, and K. D. Sales, *J. Chem. Soc. Dalton Trans.*, 185 (1973).
- (20) P. A. Cox and A. F. Orchard, *Chem. Phys. Lett.*, **7**, 273 (1970).
- (21) S. Cradock, E. A. V. Ebsworth, and A. Robertson, *J. Chem. Soc. Dalton Trans.*, 22 (1973).
- (22) For a discussion on this point, see A. H. Cowley, *Phosphorus and Sulfur*, **2**, 283 (1976).
- (23) For a review of the UPS of transition metal compounds, see A. H. Cowley, Progress in Inorganic Chemistry,
- (24) M. H. Chisholm, F. A. Cotton, and M. W. Extine, *Inorg. Chem.*, **17**, 1329 (1978).
- (25) C. E. Heath and M. B. Hursthouse, *Chem. Commun.*, 143 (1971).
- (26) R. J. Smallwood, Ph.D. Thesis, University of London, 1975.

Table I. Ionic States Resulting from Lower Energy Ionizations  
of Open-Shell Metal Dialkylamides

<u>Ground State</u>	<u>Cr(N<sup>i</sup>Pr<sub>2</sub>)<sub>3</sub></u> <u>Ion Configuration</u>	<u>Ionic State</u> <u>Produced</u>	<u>Experimental</u> <u>Ionization</u> <u>Energies<sup>a</sup></u>
$e^4 a_2^2 a_1^1 e^2 [^4A_2]$	$e^4 a_2^2 a_1^1 e^1$	$^3E$	6.53
	$e^4 a_2^2 e^2$	$^3A_2$	6.3
	$e^4 a_2^1 a_1^1 e^2$	$^3A_1$	7.38
		$^5A_1$	
	$e^3 a_2^2 a_1^1 e^2$	$^3E$	7.9
		$^5E$	
			9.9 (M-Nσ)
<u>Cr(NEt<sub>2</sub>)<sub>4</sub>, D<sub>2d</sub>(1)</u>			
$a_1^2 b_2^2 e^4 b_1^1 a_1^1 [^3B_1]$	$a_1^2 b_2^2 e^4 b_1^1$	$^2B_1$	5.9
	$a_1^2 b_2^2 e^4 a_1^1$	$^2A_1$	6.3
	$a_1^2 b_2^2 e^3 b_1^1 a_1^1$	$^2E$	7.0
		$^4E$	
	$a_1^2 b_2^1 e^4 b_1^1 a_1^1$	$^2A_2$	7.2
		$^4A_2$	
	$a_1^1 b_2^2 e^4 b_1^1 a_1^1$	$^2B_1$	7.9
		$^4B_1$	
			10.0 (M-Nσ)

Table I. (cont'd.)

<u>Cr(NEt<sub>2</sub>)<sub>4</sub>, D<sub>2d</sub>(2)</u>			
$b_1^2 e^4 a_2^2 a_1^1 b_1^1$ [ <sup>3</sup> B <sub>1</sub> ]	$b_1^2 e^4 a_2^2 a_1^1$	<sup>2</sup> A <sub>1</sub>	5.9
	$b_1^2 e^4 a_2^2 b_1^1$	<sup>2</sup> B <sub>1</sub>	6.3
	$b_1^2 e^4 a_2^1 a_1^1 b_1^1$	<sup>2</sup> B <sub>2</sub> }	7.0
		<sup>4</sup> B <sub>2</sub> }	
	$b_1^2 e^3 a_2^2 a_1^1 b_1^1$	<sup>2</sup> E }	7.2
		<sup>4</sup> E }	
	$b_1^1 e^4 a_2^2 a_1^1 b_1^1$	<sup>2</sup> A <sub>1</sub> }	7.9
		<sup>4</sup> A <sub>1</sub> }	

<sup>a</sup>All values in eV.

Table II.  $X_{\alpha}$  SW "Ground State" Eigenvalues and Charge Densities<sup>a</sup> for  $\text{Cr}(\text{NH}_2)_3$

MO	$\epsilon^b$	Cr			N			H		OUTER <sup>c</sup>		INT <sup>d</sup>	
		s	p	d	s	p	d	s		total		total	
7e	-1.00	0.03	0.63	0.66	0.01	0.06	0.07	0.00		0.02		0.11	
6e	-3.41	0.00	0.87	0.87	0.00	0.01	0.01	0.00		0.00		0.10	
4a <sub>1</sub>	-3.53	0.07	0.64	0.71	0.00	0.00	0.00	0.00		0.02		0.27	
3a <sub>2</sub>	-4.60	0.00	0.00	0.00		0.23	0.23	0.00		0.00		0.31	
5e	-5.72	0.04	0.10	0.14	0.00	0.18	0.18	0.00		0.02		0.30	
4e	-8.76	0.06	0.26	0.32	0.01	0.18	0.19	0.01		0.02		0.03	
3a <sub>1</sub>	-9.23	0.14	0.13	0.27	0.01	0.19	0.20	0.02		0.00		0.01	
2a <sub>2</sub>	-12.88	0.01		0.01		0.19	0.19	0.07		0.00		0.00	
3e	-12.93	0.00	0.02	0.02	0.00	0.19	0.19	0.07		0.01		0.00	
2e	-20.69	0.02	0.01	0.03	0.22	0.01	0.23	0.04		0.00		0.04	
2a <sub>1</sub>	-20.81	0.02	0.01	0.03	0.23	0.01	0.24	0.04		0.00		0.01	
1a <sub>2</sub>	-46.28	1.00		1.00		0.00	0.00	0.00		0.00		0.00	
1e	-46.38	0.99	0.00	0.99	0.00	0.00	0.00	0.00		0.00		0.01	
1a <sub>1</sub>	-73.51	1.00	0.00	1.00	0.00	0.00	0.00	0.00		0.00		0.00	

<sup>a</sup> charge densities for atoms are the percentages of electron densities within the atomic spheres.

<sup>b</sup> eigenvalues in eV

<sup>c</sup> charge density outside outer sphere

<sup>d</sup> intersphere charge density inside outer sphere and not accounted for by atomic spheres; dashed line denotes a separation of HOMO and LUMO

Table III. Computed Ionization Energies for  $\text{Cr}(\text{NH}_2)_3$  and  
Experimental Ionization Energies for  $\text{Cr}(\text{N}^1\text{Pr}_2)_3$ .

<u>MO</u>	<u>Computed IE<sup>a</sup></u>	<u>Experimental IE<sup>a</sup></u>
4a <sub>1</sub>	6.76	6.3
6e	7.01	6.53
3a <sub>2</sub>	7.52	7.38
5e	8.57	7.9
4e } 3a <sub>1</sub> }		9.9

<sup>a</sup>All values in eV.

Table IV. Ionization Energy Data<sup>a</sup> for  $M(NR_2)_4$  and  $M(NR_2)_5$  Compounds

<u>Ionic State</u>	<u><math>M(NR_2)_4</math></u>	
	<u><math>Mo(NMe_2)_4</math></u>	<u><math>Mo(NEt_2)_4</math></u>
${}^2B_1(d_{x^2-y^2})$	5.30	5.3
${}^2E(n_N)$	7.34	7.0 <sup>b</sup> , 7.3 <sup>b</sup>
${}^2B_2(n_N)$	7.70	7.56
${}^2A_1(n_N)$	9.01	8.7
M-N ionizations	10.7	

	<u><math>M(NR_2)_5</math></u>	
	<u><math>Nb(NMe_2)_5</math></u>	<u><math>Ta(NMe_2)_5</math></u>
$n_N$	6.77	6.89
	6.9	7.1
	7.63	7.78
	8.02	8.15
	8.21	8.35
M-N ionizations	9.7, 10.2	9.7, 10.4

<sup>a</sup>All values in eV

<sup>b</sup>Splitting due to the Jahn-Teller effect. See text.

## Figure Captions

Figure 1. He(I) UPS of : (a)  $\text{Cr}(\text{N}^i\text{Pr}_2)_3$ ; (b)  $\text{Mo}(\text{NMe}_2)_4$ ; (c)  $\text{Mo}(\text{NEt}_2)_4$ ; (d)  $\text{Cr}(\text{NEt}_2)_4$ ; and (e)  $\text{Ta}(\text{NMe}_2)_5$ .

Figure 2. Horizontal ( $\text{D}_{3h}(1)$ ,  $\phi=90^\circ$ ) and vertical ( $\text{D}_{3h}(2)$ ,  $\phi=90^\circ$ ) arrangements of N(2p) AO's in threefold symmetry.

Figure 3.  $X_\alpha$  SW contour plots for the following MO's of  $\text{Cr}(\text{NH}_2)_3$ : (a) one of the two degenerate 5e wavefunctions ( $\text{CrN}_3$  plane); (b) one of the two degenerate 4e wavefunctions ( $\text{CrN}_3$  plane); (c) the  $3a_1$  wavefunction (perpendicular to  $\text{CrN}_3$  plane, along CrN bond). Contour values are:  $1=\pm 0.20$ ;  $2=\pm 0.10$ ;  $3=\pm 0.05$ ;  $4=\pm 0.02$ .

Figure 4. A qualitative MO scheme for  $\text{M}(\text{NR}_2)_4$  complexes in two different  $\text{D}_{2d}$  configurations (see text).




Figure 1




Figure 2




Figure 3

D<sub>2d</sub> (1)

T<sub>d</sub>

D<sub>2d</sub> (2)





Figure 4

# TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. George Sandoz	1	U.S. Army Research Office P.O. Box 1211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Weapons Center China Lake, California 93555 Attn: Dr. A. B. Amster Chemistry Division	1
ONR Area Office One Hallidie Plaza, Suite 601 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Naval Civil Engineering Laboratory Port Hueneme, California 93401 Attn: Dr. R. W. Drisko	1
ONR Branch Office Building 114, Section D 666 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Professor K. E. Woehler Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R, E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1	Naval Ship Research and Development Center Annapolis, Maryland 21401 Attn: Dr. G. Bosmajian Applied Chemistry Division	1
		Naval Ocean Systems Center San Diego, California 91232 Attn: Dr. S. Yamamoto, Marine Sciences Division	1

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. R. N. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1	Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1
Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1	Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1
Dr. M. F. Hawthorne University of California Department of Chemistry Los Angeles, California 90024	1	Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 39716	1	Professor O. T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214	1
Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1	Professor P. S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1	Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1